RELIEVE PAIN/REDUCE FEVER

ByrMart

Genuine
ASPIRIN

50 TABLETS 325 mg

See carton for complete information
Active ingredient (in each tablet)
Aspirin 325 mg ..... Pain Reliever/Fever Reducer in a buffered base of Calcium Carbonate
Uses: For the temporary relief of headache, pain and fever of colds, muscle pain, menstrual pain, toothache, or minor pain of arthritis.
Warnings: Reye's syndrome: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin. **Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding. **Do not use** if you are allergic to aspirin. **Ask a doctor before use if you have** bleeding problems, asthma, ulcers, or stomach problems (such as heartburn, upset stomach, or stomach pain) that persist or recur. **Ask a doctor or pharmacist before use if you are** taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout, or arthritis. **Stop use and ask a doctor if** pain or fever persists or gets worse, redness or swelling is present, new or unexpected symptoms occur, or ringing in the ears or loss of hearing occurs. These could be signs of a serious condition. **If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.
Directions: Do not exceed recommended dosage. Do not take for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. Adults and children 12 years and over: Take 1 or 2 tablets with water every 4 to 6 hours, do not exceed 8 tablets in 24 hours unless directed by a doctor. Children under 12 years: Consult a doctor.
Other Information: Store at room temperature.
Inactive Ingredients: Carnauba Wax, Colloidal Silicon Dioxide, D&C Red#7 Calcium Lake, FD&C Blue #2 Aluminum Lake, FD&C Red #40 Aluminum Lake, Hydroxypropyl Methylcellulose, Microcrystalline Cellulose, Propylene Glycol, Shellac, Sodium Starch Glycolate, Starch, Titanium Dioxide, Zinc Stearate.
Produced especially for BYRNES PHARMACIES, GENERAL PHARMACEUTICALS, INC.
MARSHALL CITY, THOMPSON COUNTY, MARSHALL, U.S. A.

5/02
EC01

RELIEVE PAIN/REDUCE FEVER

ByrMart

Genuine
ASPIRIN

50 TABLETS 325 mg

See carton for complete information
Active ingredient (in each tablet)
Aspirin 325 mg ..... Pain Reliever/Fever Reducer in a buffered base of Calcium Carbonate
Uses: For the temporary relief of headache, pain and fever of colds, muscle pain, menstrual pain, toothache, or minor pain of arthritis.
Warnings: Reye's syndrome: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin. **Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding. **Do not use** if you are allergic to aspirin. **Ask a doctor before use if you have** bleeding problems, asthma, ulcers, or stomach problems (such as heartburn, upset stomach, or stomach pain) that persist or recur. **Ask a doctor or pharmacist before use if you are** taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout, or arthritis. **Stop use and ask a doctor if** pain or fever persists or gets worse, redness or swelling is present, new or unexpected symptoms occur, or ringing in the ears or loss of hearing occurs. These could be signs of a serious condition. **If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.
Directions: Do not exceed recommended dosage. Do not take for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. Adults and children 12 years and over: Take 1 or 2 tablets with water every 4 to 6 hours, do not exceed 8 tablets in 24 hours unless directed by a doctor. Children under 12 years: Consult a doctor.
Other Information: Store at room temperature.
Inactive Ingredients: Carnauba Wax, Colloidal Silicon Dioxide, D&C Red#7 Calcium Lake, FD&C Blue #2 Aluminum Lake, FD&C Red #40 Aluminum Lake, Hydroxypropyl Methylcellulose, Microcrystalline Cellulose, Propylene Glycol, Shellac, Sodium Starch Glycolate, Starch, Titanium Dioxide, Zinc Stearate.
Produced especially for BYRNES PHARMACIES, GENERAL PHARMACEUTICALS, INC.
MARSHALL CITY, THOMPSON COUNTY, MARSHALL, U.S. A.

5/02
EC01

RELIEVE PAIN/REDUCE FEVER

ByrMart

Genuine
ASPIRIN

50 TABLETS 325 mg

See carton for complete information
Active ingredient (in each tablet)
Aspirin 325 mg ..... Pain Reliever/Fever Reducer in a buffered base of Calcium Carbonate
Uses: For the temporary relief of headache, pain and fever of colds, muscle pain, menstrual pain, toothache, or minor pain of arthritis.
Warnings: Reye's syndrome: Children and teenagers should not use this medicine for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin. **Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding. **Do not use** if you are allergic to aspirin. **Ask a doctor before use if you have** bleeding problems, asthma, ulcers, or stomach problems (such as heartburn, upset stomach, or stomach pain) that persist or recur. **Ask a doctor or pharmacist before use if you are** taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout, or arthritis. **Stop use and ask a doctor if** pain or fever persists or gets worse, redness or swelling is present, new or unexpected symptoms occur, or ringing in the ears or loss of hearing occurs. These could be signs of a serious condition. **If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.
Directions: Do not exceed recommended dosage. Do not take for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. Adults and children 12 years and over: Take 1 or 2 tablets with water every 4 to 6 hours, do not exceed 8 tablets in 24 hours unless directed by a doctor. Children under 12 years: Consult a doctor.
Other Information: Store at room temperature.
Inactive Ingredients: Carnauba Wax, Colloidal Silicon Dioxide, D&C Red#7 Calcium Lake, FD&C Blue #2 Aluminum Lake, FD&C Red #40 Aluminum Lake, Hydroxypropyl Methylcellulose, Microcrystalline Cellulose, Propylene Glycol, Shellac, Sodium Starch Glycolate, Starch, Titanium Dioxide, Zinc Stearate.
Produced especially for BYRNES PHARMACIES, GENERAL PHARMACEUTICALS, INC.
MARSHALL CITY, THOMPSON COUNTY, MARSHALL, U.S. A.

5/02
EC01

Labels of ByrMart aspirin bottles
Recovered at 131 N. Lombard, 12 September 2001